

Reg. No. 10-B-2-56

# COPPER OXYCHLORIDE 85WP

FUNGICIDE

A fungicide for the control of certain diseases in agricultural and horticultural crops, in flowers and ornamentals.

<b>Composition</b>	<b>mass/mass</b>
Copper oxychloride	85 %
Inert ingredients	15 %
Chemical group:	Inorganic Compound

20 litres



**POISON**

**TO CAUSE A HAZARD IN THE USE, STORAGE OR DISPOSAL OF THIS SUBSTANCE IS AN OFFENCE.**

**NOVA-AGRO**  
CHEMICALS

4 Hood Rd | Southerton | Harare

**DIRECTIONS FOR USE:**

Use only as directed

Ensure adequate creaming before dilution. Agitate the spray mixture constantly whilst spraying. Apply when the foliage is dry as a full cover spray so as to wet plants thoroughly.

Crop	Disease	Mixture of Copper Oxychloride in water	Application	Min. days from last spray to harvest
Citrus	Black spot	200g/100 litre	Apply at 75% petal drop as a cover spray. Repeat twice at 4-5 week intervals.	14
Coffee	Rust	5-8kg/ha	Apply in sufficient water to give thorough coverage.	14
Potatoes	Early and Late blight	500g/100 litre	Use at the rate of 550 to 1100 litre per hectare, commencing when the potatoes emerge. Repeat every 7 days or after every 40mm of rain.	14
Tobacco Seedbeds (Always consult the TRB Handbook)	Wildlife Angular leaf spot		Apply weekly. Start spraying four weeks after emergence.	N.a.
		300g-500g/100 litres	High volume spray: Apply the sprays at 200ml/m <sup>2</sup> and increase as the seedlings grow to a max. of 500ml/m <sup>2</sup> . If either disease is seen, apply the higher dosage in 500ml/m <sup>2</sup> , to give complete coverage.	
		100g-465g/litre	Ultra low volume spray: Apply at 4,0ml/m <sup>2</sup> and if disease is seen, increase the dosage rate.	
Tomatoes	Early and late blight	500g/100 litre	Apply as overall spray before disease appears. Repeat at 7-day intervals or in the summer after every 40mm of rain, whichever is the soonest.	3

**COMPATIBILITY**

The manufacturers stress that it is preferable to use this product on its own; furthermore, the quality of water in a mixture may adversely affect its compatibility. The suggestions below are therefore guidelines only, and all responsibility is disclaimed for lack of effectiveness, damage or loss of any kind from mixing this product with another. The manufacturers strongly recommend a trial mixture be taken with water intended for use in the main mixture. Subject to the above, this product may normally be mixed at those water concentrations common in the use of conventional ground spraying equipment with any ONE of the following: Mancozeb 80WP, Dimethoate EC, Carbaryl 85WP, Endosulfan 35EC and Endosulfan 50WP.

In tobacco however, this product should not be mixed with any other fungicides used in seedbeds.

**PRECAUTIONS:**

1. Handle with care and avoid any contact with the pesticide.
2. WEAR PROTECTIVE CLOTHING i.e. gloves and overalls.
3. Do not eat, drink or smoke while applying this pesticide.
4. DANGEROUS TO ANIMALS, BIRDS, FISH AND BEES.
5. KEEP OUT OF REACH OF CHILDREN AND AWAY FROM FOODS AND STORE UNDER LOCK AND KEY.
6. Destroy empty container – DO NOT use for any other purpose.

**FIRST AID:**

Should accidental poisoning occur call a doctor and show him this label. In case of contact with: - Eyes: Wash eyes with clean water for at least 15 minutes. Skin: Remove contaminated clothing. Wash skin with plenty of soap and water. If swallowed, induce vomiting. If inhaled, remove patient to fresh air and loosen clothing around the throat and chest.

**ANTIDOTE:**

No known antidote.

**WARRANTY:**

The excellent results which this chemical should normally yield are not however warranted or guaranteed, as effectiveness can be greatly influenced by factors outside the control of either the manufacturers or the suppliers. No warranty whatsoever, express or implied, is therefore given concerning the performance or effectiveness of this chemical and responsibility is specifically excluded for any damage, injury or loss of any kind during or resulting from its improper handling, use or storage, whether or not such has been in accordance with directions.

**DATE OF MANUFACTURE:****BATCH NO:**

Manufactured by:

NOVA AGRO (HK) LTD  
28/F Tesbury Centre  
28 Queen's Road East  
Wanchai, HONG KONG

Registration held by/distributed by:

**NOVA-AGRO**  
**CHEMICALS**

4 Hood Road | Southerton | Harare