

#### SAFETY PRECAUTIONS

1. Handle with care. Avoid contact/splashing.
2. Wear protective clothing - wear rubber gloves and face shield when handling the concentrate.
3. KEEP OUT OF THE REACH OF CHILDREN.
4. KEEP AWAY FROM FOOD AND FOODSTUFFS.
5. KEEP UNDER LOCK AND KEY.
6. DO NOT eat, drink or smoke whilst handling this product.
7. Invert the empty container over the spray or mixing tank and allow to drain for at least 30 seconds after the flow has slowed down to a drip. Thereafter, rinse the container three times with a volume of water equal to a minimum of 10% of that container. Add the rinsings to the contents of the spray tank before destroying the container.
8. While spraying, avoid contact with the spray as much as you can since product may cause eye and skin irritation. Avoid spray drift onto other crops, grazing, rivers or dams.
9. Clean applicator thoroughly after use and dispose of wash water where it will not contaminate crops, grazing, rivers or dams. Toxic to fish and harmful to other aquatic life.
10. Destroy empty container and do not use for any other purpose.
11. Change and wash your work clothes after use. Wash yourself.

#### SYMPTOMS OF POISONING

Symptoms similar to those caused by nicotine poisoning. Apathetic state, depressed muscular tone, respiratory disturbances and trembling; muscular cramps also possible in severe cases of poisoning.

#### FIRST AID

1. If any of the above symptoms occur, particularly if contamination is likely, STOP WORK.
2. Remove contaminated clothing and wash skin and hair thoroughly.
3. If swallowed, induce vomiting without delay by tickling the back of the throat.
4. CALL A DOCTOR AND SHOW HIM THIS LABEL.

#### NOTE TO PHYSICIAN

No specific antidote. Treat symptomatically.

#### WARNINGS

1. Allow 32 days between last application and harvest of maize.
2. On tobacco, IMIDACLOPRID 200SL should only be used at planting and should not be applied over the crown of the seedling.

#### WARRANTY

Provided that the recommendations for usage of this product are strictly followed it should give excellent results, but since the performance of crop protection chemicals and pesticides can be influenced by factors outside the manufacturer's control, the manufacturers and seller give no guarantee or warranty, either express or implied, concerning the use and/or storage of this product, and accept no responsibility or liability of any nature whatsoever for any damage, injury or financial loss of any nature from the use thereof.

Manufactured by:

NOVAAGRO (HK) LTD  
6TH FLOOR, WYNDHAM PLACE,  
44 WYNDHAM STREET,  
CENTRAL HONG KONG


# IMIDACLOPRID 200SL

Insecticide/Termicide

Reg. No.: 10-D-142-41

TRB. No.: 16-19-D-48

1L



HARMFUL IF SWALLOWED

**Composition mass/vol.**  
Imidacloprid.....200g/L  
Inert ingredients.....to 1 litre

**Chemical group:** neonicotinoid

#### INSECTICIDE/TERMITICIDE FOR AGRICULTURAL USE

A systemic insecticide for the control of aphids and termites in tobacco as well as termites in maize and cotton.

**TO CAUSE A HAZARD IN THE USE, STORAGE OR DISPOSAL OF THIS SUBSTANCE IS AN OFFENCE.**

#### DIRECTIONS FOR USE - Use only as directed.

Pour the required quantity of IMIDACLOPRID 200SL into the water while agitating thoroughly. Do not allow the IMIDACLOPRID 200SL mixture to stand any length of time before using.

#### APPLICATION RATES

CROP	PEST	DOSAGE RATE/1000	REMARKS
Tobacco lands TRB No. 16-19-D-48 (Always consult TRB Handbook for detailed recommendations)	Aphids	1,0L/ha at planting	Apply at an overall rate of 1,0L/ha at planting using one of the following methods: <b>Mix directly in bulk planting water supply:</b> Pour 1L of mix into each planting hole. <b>Mix separately from planting water:</b> Apply one 30ml cup of mix in each planting hole immediately after filling it prior to planting.
	Termites	6,5ml/100L water 220ml/100L water	
Maize	Termites	200ml (0,5L/ha)	Apply 8 - 10 weeks after planting (prior to termite attack). Apply, using a knapsack, as a 30cm band in the maize row with at least 250L water/ha.
Cotton	Termites	1,0L/ha	<b>Furrow treatment:</b> Apply over seed in the furrow before covering with soil. Use a knapsack applying at least 100L spray mixture/ha.

#### COMPATIBILITY

IMIDACLOPRID 200SL is compatible with most conventional insecticides.

#### SAFETY PRECAUTIONS

##### Empty container disposal

Invert the empty container over the spray tank or mixing tank and allow to drain for at least 30 seconds after the flow has slowed down to a drip. Thereafter, rinse the container three times with a volume of water equal to a minimum of 10% of the container. Add the rinsings to the contents of the spray tank. Destroy the empty container by perforation and flattening. Return to supplier for recycling. DO NOT use for any other purpose. Dispose of the wash water at a site for the disposal of pesticides.

##### Decontamination of sprayer

Clean applicator thoroughly after use and ensure that all traces of IMIDACLOPRID 200SL are removed. Make use of the following method:

- (a) Drain and rinse tank, spray boom and hoses with clean water for at least 10 minutes
- (b) Fill tank with clean water and add it to 1 litre household bleach (5%) or 1.5 Litres of household bleach (3.5%) per 200 Litres of water. Rinse hoses and spray boom and leave in tank for 15 minutes whilst agitating. Drain through nozzle outlets.
- (c) Repeat step (b) and thereafter, rinse thoroughly with clean water and dispose of the wash water at a site designated for the disposal of pesticides.

Date of Manufacture:

Batch No.:

Registration held by:

MAGCHEM (PVT) LIMITED  
2274 TILBURY ROAD,  
WORKINGTON, HARARE,  
ZIMBABWE